

THE FOURTEENTH DATA RELEASE OF THE SLOAN DIGITAL SKY SURVEY: FIRST SPECTROSCOPIC DATA FROM THE EXTENDED BARYON OSCILLATION SKY SURVEY AND FROM THE SECOND PHASE OF THE APACHE POINT OBSERVATORY GALACTIC EVOLUTION EXPERIMENT

BELA ABOLFATHI¹, D. S. AGUADO², GABRIELA AGUILAR³, CARLOS ALLENDE PRIETO^{2,4}, ANDRES ALMEIDA⁵, TONIMA TASNIM ANANNA⁶, FRIEDRICH ANDERS⁷, SCOTT F. ANDERSON⁸, BRETT H. ANDREWS⁹, BORJA ANGUIANO¹⁰, ALFONSO ARAGÓN-SALAMANCA¹¹, MARIA ARGUDO-FERNÁNDEZ¹², ERIC ARMENGAUD¹³, METIN ATA⁷, ERIC AUBOURG¹⁴, VLADIMIR AVILA-REESE³, CARLES BADENES⁹, STEPHEN BAILEY¹⁵, KATHLEEN A. BARGER¹⁶, JORGE BARRERA-BALLESTEROS¹⁷, CURTIS BARTOSZ⁸, DOMINIC BATES¹⁸, FALK BAUMGARTEN^{7,19}, JULIAN BAUTISTA²⁰, RACHAEL BEATON²¹, TIMOTHY C. BEERS²², FRANCESCO BELFIORE^{23,24,25}, CHAD F. BENDER²⁶, MARIANGELA BERNARDI²⁷, MATTHEW A. BERSHADY²⁸, FLORIAN BEUTLER²⁹, JONATHAN C. BIRD³⁰, DMITRY BIZYAEV^{31,32,33}, GUILLERMO A. BLANC²¹, MICHAEL R. BLANTON³⁴, MICHAEL BLOMQUIST³⁵, ADAM S. BOLTON³⁶, MÉDÉRIC BOQUIEN¹², JURA BORISSOVA^{37,38}, JO BOVY^{39,40,41}, CHRISTIAN ANDRES BRADNA DIAZ⁴², WILLIAM NIELSEN BRANDT^{43,44,45}, JONATHAN BRINKMANN³¹, JOEL R. BROWNSTEIN²⁰, KEVIN BUNDY²⁵, ADAM J. BURGASSER⁴⁶, ETIENNE BURTIN¹³, NICOLÁS G. BUSCA¹⁴, CALEB I. CAÑAS⁴³, MARIANA CANO-DÍAZ⁴⁷, MICHELE CAPPELLARI⁴⁸, RICARDO CARRERA^{2,4}, ANDREW R. CASEY⁴⁹, YANPING CHEN⁵⁰, BRIAN CHERINKA⁵¹, CRISTINA CHIAPPINI⁷, PETER DOOHYUN CHOI⁵², DREW CHOJNOWSKI³², CHIA-HSUN CHUANG⁷, HAEUN CHUNG⁵³, NICOLAS CLERC^{54,55,56}, ROGER E. COHEN^{57,58}, JOHAN COMPARAT⁵⁴, JANAINA CORREA DO NASCIMENTO^{59,60}, LUIZ DA COSTA^{60,61}, MARIE-CLAUDE COUSINOU⁶², KEVIN COVEY⁶³, JEFFREY D. CRANE²¹, IRENE CRUZ-GONZALEZ³, KATIA CUNHA^{61,26}, GUILLERMO J. DAMKE^{10,64,65}, JEREMY DARLING⁶⁶, JAMES W. DAVIDSON JR.¹⁰, KYLE DAWSON²⁰, ANNA BÁRBARA DE ANDRADE QUEIROZ^{59,60}, MIGUEL ANGEL C. DE ICАЗA LIZAOLA³, AXEL DE LA MACORRA⁶⁷, SYLVAIN DE LA TORRE³⁵, NATHAN DE LEE^{68,30}, VICTORIA DE SAINTE AGATHE⁶⁹, ALICE DECONTO MACHADO^{70,60}, FLAVIA DELL'AGLI^{2,4}, TIMOTHÉE DELUBAC⁷¹, ALEKSANDAR M. DIAMOND-STANIC⁴², JOHN DONOR¹⁶, JUAN JOSÉ DOWNES⁷², NIV DRORY⁷³, HÉLION DU MAS DES BOURBOUX¹³, CHRISTOPHER J. DUCKWORTH¹⁸, TOM DWELLY⁵⁴, JAMIE DYER²⁰, GARRETT EBELKE¹⁰, ARTHUR DAVIS EIGENBROT²⁸, DANIEL J. EISENSTEIN⁷⁴, YVONNE P. ELSWORTH⁷⁵, ERIC EMMELLEN⁷⁶, MIKE ERACLEOUS⁴⁴, STEPHANIE ESCOFFIER⁶², XIAOHUI FAN²⁶, EMMA FERNÁNDEZ ALVAR³, J. G. FERNÁNDEZ-TRINCADO⁵⁷, RAFAEL FERNANDO CIROLINI⁶⁰, DIANE FEUILLET⁷⁷, SCOTT W. FLEMING⁵⁸, ANDREU FONT-RIBERA⁷⁸, GORDON FREISCHLAD³¹, PETER FRINCHABOY¹⁶, YILEN GÓMEZ MAQUEO CHEW³, LLUÍS GALBANY⁹, ANA E. GARCÍA PÉREZ², R. GARCIA-DIAS^{2,4}, D. A. GARCÍA-HERNÁNDEZ^{2,4}, PATRICK GAULME³¹, JOSEPH GELFAND³⁴, HÉCTOR GIL-MARÍN^{79,80}, BRUCE A. GILLESPIE³¹, DANIEL GODDARD²⁹, VIOLETA GONZALEZ-PEREZ²⁹, KATHLEEN GRABOWSKI³¹, PAUL J. GREEN⁷⁴, CATHERINE J. GRIER^{43,44}, ALAIN GUEGUEN⁵⁴, HONG GUO⁸¹, JULIEN GUY⁶⁹, ALEX HAGEN⁴⁴, PAUL HARDING⁸², STEN HASSELQUIST³², SUZANNE HAWLEY⁸, FRED HEARTY⁴³, SASKIA HEKKER⁸³, JESUS HERNÁNDEZ³, HECTOR HERNANDEZ TOLEDO³, DAVID W. HOGG³⁴, KELLY HOLLEY-BOCKELMANN³⁰, JON HOLTZMAN³², JIAMIN HOU⁵⁴, BAU-CHING HSIEH⁸⁴, JASON A. S. HUNT⁴⁰, TIMOTHY A. HUTCHINSON²⁰, HO SEONG HWANG⁵³, CAMILO EDUARDO JIMENEZ ANGEL^{2,4}, JENNIFER A. JOHNSON^{85,86}, AMY JONES⁸⁷, HENRIK JÖNSSON^{2,4}, ERIC JULLO³⁵, FAHIM SAKIL KHAN⁴², KAREN KINEMUCHI³¹, DAVID KIRKBY¹, CHARLES C. KIRKPATRICK IV⁸⁸, FRANCISCO-SHU KITAUURA^{2,4}, GILLIAN R. KNAPP⁸⁹, JEAN-PAUL KNEIB⁷¹, JUNA A. KOLLMEIER²¹, IVAN LACERNA^{38,90,91}, RICHARD R. LANE⁹⁰, DUSTIN LANG⁴⁰, DAVID R. LAW⁵⁸, JEAN-MARC LE GOFF¹³, YOUNG-BAE LEE⁵², HONGYU LI⁹², JIANHUI LIAN²⁹, YU LIANG⁹³, LIHWAI LIN (林俐暉)⁸⁴, DAN LONG³¹, SARA LUCATELLO⁹⁴, BRITT LUNDGREN⁹⁵, J. TED MACKERETH⁹⁶, CHELSEA L. MACLEOD⁷⁴, SUVRATH MAHADEVAN⁴³, MARCIO ANTONIO-GEIMBA MAIA^{61,60}, STEVEN MAJEWSKI¹⁰, ARTURO MANCHADO^{2,4}, CLAUDIA MARASTON²⁹, VIVEK MARIAPPAN²⁰, RUI MARQUES-CHAVES^{2,4}, THOMAS MASSERON^{2,4}, KAREN L. MASTERS (何凱諭)^{29,97}, RICHARD M. McDERMID⁹⁸, IAN D. MCGREER²⁶, MATTHEW MELENDEZ¹⁶, SOFIA MENESES-GOYTIA²⁹, ANDREA MERLONI⁵⁴, MICHAEL R. MERRIFIELD¹¹, SZABOLCS MESZAROS^{99,100}, ANDRES MEZA¹⁰¹, IVAN MINCHEV⁷, DANTE MINNITI^{101,38,102}, EVA-MARIA MUELLER²⁹, FRANCISCO MULLER-SANCHEZ⁶⁶, DEMITRI MUNA⁸⁶, ADAM D. MYERS¹⁰³, PREETHI NAIR¹⁰⁴, KIRPAL NANDRA⁵⁴, MELISSA NESS⁷⁷, JEFFREY A. NEWMAN⁹, ROBERT C. NICHOL²⁹, DAVID L. NIDEVER³⁶, CHRISTIAN NITSCHHELM¹², PASQUIER NOTERDAEME¹⁰⁵, JULIA O'CONNELL¹⁶, RYAN JAMES OELKERS³⁰, AUDREY ORAVETZ³¹, DANIEL ORAVETZ³¹, ERIK AQUINO ORTÍZ³, YEISSON OSORIO^{2,4}, ZACH PACE²⁸, NELSON PADILLA⁹⁰, NATHALIE PALANQUE-DELABROUILLE¹³, PEDRO ALONSO PALICIO^{2,4}, HSI-AN PAN⁸⁴, KAIKE PAN³¹, TANIYA PARIKH²⁹, ISABELLE PÂRIS³⁵, CHANGBOM PARK⁵³, SEBASTIEN PEIRANI¹⁰⁵, MARCOS PELLEJERO-IBANEZ^{2,4}, SAMANTHA PENNY²⁰, WILL J. PERCIVAL²⁹, ISMAEL PEREZ-FOURNON^{2,4}, PATRICK PETITJEAN¹⁰⁵, MATTHEW M. PIERI³⁵, MARC PINSONNEAULT⁸⁵, ALICE PISANI⁶², FRANCISCO PRADA^{106,107}, ABHISHEK PRAKASH⁹, M. JORDAN RADDICK¹⁷, ANAND RAICHOOR⁷¹, SANDRO BARBOZA REMBOLD^{70,60}, HANNAH RICHSTEIN¹⁶, ROGEMAR A. RIFFEL^{70,60}, ROGÉRIO RIFFEL^{59,60}, HANS-WALTER RIX⁷⁷, ANNIE C. ROBIN¹⁰⁸, SERGIO RODRÍGUEZ TORRES¹⁰⁹, CARLOS ROMÁN-ZÚNIGA¹¹⁰, ASHLEY J. ROSS⁸⁶, GRAZIANO ROSSI⁵², JOHN RUAN⁸, ROSSANA RUGGERI²⁹, JOSE RUIZ⁴², MARA SALVATO⁵⁴, ARIEL G. SÁNCHEZ⁵⁴, SEBASTIÁN F. SÁNCHEZ³, JORGE SANCHEZ ALMEIDA², JOSÉ R. SÁNCHEZ-GALLEGO⁸, BASÍLIO XAVIER SANTIAGO^{59,60}, RICARDO P. SCHIAVON⁹⁶, JADERSON S. SCHIMOIA^{59,60}, EDWARD SCHLAFLY¹⁵, DAVID SCHLEGEL¹⁵, DONALD P. SCHNEIDER^{43,44}, WILLIAM J. SCHUSTER^{3,110}, AXEL SCHWOPE⁷, HEE-JONG SEO¹¹¹, ALDO SERENELLI¹¹², SHIYIN SHEN⁸¹, YUE SHEN^{113,114}, MATTHEW SHETRONE⁷³, MICHAEL SHULL⁶⁶, VÍCTOR SILVA AGUIRRE¹¹⁵, JOSHUA D. SIMON²¹, MIKE SKRUTSKIE¹⁰, ANŽE SLOSAR¹¹⁶, REBECCA SMETHURST¹¹, VERNE SMITH³⁶, JENNIFER SOBECK⁸, GARRETT SOMERS³⁰, BONNIE J. SOUTER¹⁷, DIOGO SOUTO⁶¹, ASHLEY SPINDLER¹¹⁷, DAVID V. STARK¹¹⁸, KEIVAN STASSUN³⁰, MATTHIAS STEINMETZ⁷, DENNIS STELLO^{119,120,115}, THAISA STORCHI-BERGMANN^{59,60}, ALINA STREBLYANSKA^{2,4}, GUY STRINGFELLOW⁶⁶, GENARO SUÁREZ³, JING SUN¹⁶, LASZLO SZIGETI⁹⁹, MANUCHEHR TAGHIZADEH-POPP¹⁷, MICHAEL S. TALBOT²⁰, BAITIAN TANG⁵⁷, CHARLING TAO^{93,62}, JAMIE TAYAR⁸⁵, MITA TEMBE¹⁰, DANIEL THOMAS²⁹, PATRICIA TISSERA¹⁰¹, RITA TOJEIRO¹⁸, CHRISTY TREMONTI²⁸, NICHOLAS W. TROUP¹⁰, O. VALENZUELA⁴⁷, REMCO VAN DEN BOSCH⁷⁷, JAIME VARGAS-GONZÁLEZ⁶⁵, MARIANA VARGAS-MAGAÑA⁶⁷, JOSE ALBERTO VAZQUEZ¹¹⁶, SANDRO VILLANOVA⁵⁷, NICOLE VOGT³², DAVID WAKE^{117,95}, YUTING WANG⁹², BENJAMIN ALAN WEAVER³⁶, ANNE-MARIE WEIJMANS¹⁸, KYLE B. WESTFALL¹²¹, DAVID G. WHELAN¹²², ERIC WILCOTS²⁸, VIVIENNE WILD¹⁸, ROB A. WILLIAMS⁹⁶, JOHN WILSON¹⁰,

W. M. WOOD-VASEY⁹, DOMINIKA WYLEZALEK⁵¹, TING XIAO (肖婷)⁸¹, RENBIN YAN¹²³, MENG YANG¹⁸, JASON E. YBARRA¹²⁴, CHRISTOPHE YÈCHE¹³, NADIA ZAKAMSKA¹⁷, OLGA ZAMORA^{2,4}, PAULINE ZARROUK¹³, GAIL ZASOWSKI^{58,20}, KAI ZHANG¹²³, CHENG ZHAO⁹³, GONG-BO ZHAO^{92,29}, ZHENG ZHENG²⁰, ZHI-MIN ZHOU⁹², GUANGTUN ZHU^{51,125}, JOEL C. ZINN⁸⁵, HU ZOU⁹²

Draft version July 28, 2017

Abstract

The fourth generation of the Sloan Digital Sky Survey (SDSS-IV) has been in operation since July 2014. This paper describes the second data release from this phase, and the fourteenth from SDSS overall (making this, Data Release Fourteen or DR14). This release makes public data taken by SDSS-IV in its first two years of operation (July 2014–2016). Like all previous SDSS releases, DR14 is cumulative, including the most recent reductions and calibrations of all data taken by SDSS since the first phase began operations in 2000.

New in DR14 is the first public release of data from the extended Baryon Oscillation Sky Survey (eBOSS); the first data from the second phase of the Apache Point Observatory (APO) Galactic Evolution Experiment (APOGEE-2), including stellar parameter estimates from an innovative data driven machine learning algorithm known as The Cannon; and almost twice as many data cubes from the Mapping Nearby Galaxies at APO (MaNGA) survey as were in the previous release ($N = 2812$ in total).

This paper describes the location and format of the publicly available data from SDSS-IV surveys. We provide references to the important technical papers describing how these data have been taken (both targeting and observation details) and processed for scientific use. The SDSS website (www.sdss.org) has been updated for this release, and provides links to data downloads, as well as tutorials and examples of data use.

SDSS-IV is planning to continue to collect astronomical data until 2020.

Subject headings: Atlases — Catalogs — Surveys

spokesperson@sdss.org

¹ Department of Physics and Astronomy, University of California, Irvine, Irvine, CA 92697, USA

² Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

³ Instituto de Astronomía, Universidad Nacional Autónoma de México, A.P. 70-264, 04510, México, D.F., México

⁴ Departamento de Astrofísica, Universidad de La Laguna (ULL), E-38206 La Laguna, Tenerife, Spain

⁵ Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, Universidad de La Serena, Benavente 980, La Serena, Chile

⁶ Yale Center for Astronomy and Astrophysics, Yale University, New Haven, CT, 06520, USA

⁷ Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, D-14482 Potsdam, Germany

⁸ Department of Astronomy, Box 351580, University of Washington, Seattle, WA 98195, USA

⁹ PITT PACC, Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA 15260, USA

¹⁰ Department of Astronomy, University of Virginia, 530 McCormick Road, Charlottesville, VA 22904-4325, USA

¹¹ School of Physics & Astronomy, University of Nottingham, Nottingham, NG7 2RD, United Kingdom

¹² Unidad de Astronomía, Fac. Cs. Básicas, Universidad de Antofagasta, Avda. U. de Antofagasta 02800, Antofagasta, Chile

¹³ CEA, Centre de Saclay, IRFU, F-91191, Gif-sur-Yvette, France

¹⁴ APC, University of Paris Diderot, CNRS/IN2P3, CEA/IRFU, Observatoire de Paris, Sorbonne Paris Cité, France

¹⁵ Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA

¹⁶ Department of Physics and Astronomy, Texas Christian University, Fort Worth, TX 76129, USA

¹⁷ Department of Physics and Astronomy, Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218, USA

¹⁸ School of Physics and Astronomy, University of St Andrews, North Haugh, St Andrews, KY16 9SS

¹⁹ Humboldt-Universität zu Berlin, Institut für Physik, Newtonstrasse 15, D-12589, Berlin, Germany

²⁰ Department of Physics and Astronomy, University of Utah,

115 S. 1400 E., Salt Lake City, UT 84112, USA

²¹ The Observatories of the Carnegie Institution for Science, 813 Santa Barbara St., Pasadena, CA 91101, USA

²² Department of Physics and JINA Center for the Evolution of the Elements, University of Notre Dame, Notre Dame, IN 46556, USA

²³ Cavendish Laboratory, University of Cambridge, 19 J. J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom

²⁴ Kavli Institute for Cosmology, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

²⁵ University of California Observatories, University of California, Santa Cruz, CA 95064, USA

²⁶ Steward Observatory, The University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721-0065, USA

²⁷ Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, USA

²⁸ Department of Astronomy, University of Wisconsin-Madison, 475 N. Charter St., Madison, WI 53726, USA

²⁹ Institute of Cosmology & Gravitation, University of Portsmouth, Dennis Sciama Building, Portsmouth, PO1 3FX, UK

³⁰ Vanderbilt University, Department of Physics & Astronomy, 6301 Stevenson Center Ln., Nashville, TN 37235, USA

³¹ Apache Point Observatory, P.O. Box 59, Sunspot, NM 88349, USA

³² Department of Astronomy, New Mexico State University, Box 30001, MSC 4500, Las Cruces NM 88003, USA

³³ Sternberg Astronomical Institute, Moscow State University, Moscow

³⁴ Center for Cosmology and Particle Physics, Department of Physics, New York University, 726 Broadway, Room 1005, New York, NY 10003, USA

³⁵ Aix Marseille Univ, CNRS, LAM, Laboratoire d'Astrophysique de Marseille, Marseille, France

³⁶ National Optical Astronomy Observatory, 950 North Cherry Avenue, Tucson, AZ 85719, USA

³⁷ Departamento de Física y Astronomía, Universidad de Valparaíso, Av. Gran

³⁸ Instituto Milenio de Astrofísica, Av. Vicuña Mackenna 4860, Macul, Santiago, Chile

³⁹ Department of Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON, M5S 3H4, Canada

⁴⁰ Dunlap Institute for Astronomy and Astrophysics, Univer-

1. INTRODUCTION

It is now sixteen years since the first data release from the Sloan Digital Sky Survey (SDSS; York et al. 2000). This Early Data Release, or EDR, occurred in June 2001 (Stoughton et al. 2002). Since this time, annual data releases from SDSS have become part of the landscape of astronomy, making the SDSS's 2.5 meter Sloan Foundation Telescope (Gunn et al. 2006) one of the most pro-

ductive observatories in the world (Madrid & Macchetto 2009), and populating databases used by thousands of astronomers worldwide (Raddick et al. 2014a,b). This paper describes the fourteenth public data release from SDSS, or DR14, released on 31st July 2017.

The SDSS has completed three phases and is currently in its fourth phase. SDSS-I and -II conducted a Legacy survey of galaxies and quasars (York et al. 2000), the

sity of Toronto, 50 St. George Street, Toronto, Ontario M5S 3H4, Canada

⁴¹ Alfred P. Sloan Fellow

⁴² Department of Physics and Astronomy, Bates College, 44 Campus Avenue, Lewiston, ME 04240, USA

⁴³ Department of Astronomy and Astrophysics, Eberly College of Science, The Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802, USA

⁴⁴ Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA

⁴⁵ Department of Physics, The Pennsylvania State University, University Park, PA 16802, USA

⁴⁶ Center for Astrophysics and Space Science, University of California San Diego, La Jolla, CA 92093, USA

⁴⁷ CONACYT Research Fellow, Instituto de Astronomía, Universidad Nacional Autónoma de México, A.P. 70-264, 04510, México, D.F., México

⁴⁸ Sub-department of Astrophysics, Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

⁴⁹ School of Physics & Astronomy, Monash University, Wellington Road, Clayton, Victoria 3800, Australia

⁵⁰ NYU Abu Dhabi, PO Box 129188, Abu Dhabi, UAE

⁵¹ Center for Astrophysical Sciences, Department of Physics and Astronomy, Johns Hopkins University, 3400 North Charles Street, Baltimore, MD 21218, USA

⁵² Department of Astronomy and Space Science, Sejong University, Seoul 143-747, Korea

⁵³ Korea Institute for Advanced Study, 85 Hoegiro, Dongdaemun-gu, Seoul 02455, Republic of Korea

⁵⁴ Max-Planck-Institut für Extraterrestrische Physik, Gießenbachstr. 1, D-85748 Garching, Germany

⁵⁵ CNRS, IRAP, 9 Av.Colonel Roche, BP 44346, F-31028 Toulouse cedex 4, France

⁵⁶ Université de Toulouse, UPS-OMP, IRAP, Toulouse, France

⁵⁷ Departamento de Astronomia, Casilla 160-C, Universidad de Concepcion, Concepcion, Chile

⁵⁸ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁵⁹ Instituto de Física, Universidade Federal do Rio Grande do Sul, Campus do Vale, Porto Alegre, RS, Brasil, 91501-970

⁶⁰ Laboratório Interinstitucional de e-Astronomia, 77 Rua General José Cristino, Rio de Janeiro, 20921-400, Brasil

⁶¹ Observatório Nacional, Rio de Janeiro, Brasil

⁶² Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

⁶³ Department of Physics and Astronomy, Western Washington University, 516 High Street, Bellingham, WA 98225, USA

⁶⁴ Centro Multidisciplinario de Ciencia y Tecnología, Universidad de La Serena, Cisternas 1200, La Serena, Chile

⁶⁵ Departamento de Física, Facultad de Ciencias, Universidad de La Serena, Cisternas 1200, La Serena, Chile

⁶⁶ Center for Astrophysics and Space Astronomy, Department of Astrophysical and Planetary Sciences, University of Colorado, 389 UCB, Boulder, CO 80309-0389, USA

⁶⁷ Instituto de Física, Universidad Nacional Autónoma de México, Apdo. Postal 20-364, México.

⁶⁸ Department of Physics, Geology, and Engineering Tech, Northern Kentucky University, Highland Heights, KY 41099, USA

⁶⁹ LPNHE, CNRS/IN2P3, Université Pierre et Marie Curie Paris 6, Université Denis Diderot Paris, 4 place Jussieu, 75252 Paris CEDEX, France

⁷⁰ Departamento de Física, CCNE, Universidade Federal de Santa Maria, 97105-900, Santa Maria, RS, Brazil

⁷¹ Institute of Physics, Laboratory of Astrophysics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Observatoire de

Sauverny, 1290 Versoix, Switzerland

⁷² Centro de Investigaciones de Astronomía, AP 264, Mérida 5101-A, Venezuela

⁷³ McDonald Observatory, The University of Texas at Austin, 1 University Station, Austin, TX 78712, USA

⁷⁴ Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA

⁷⁵ School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

⁷⁶ European Southern Observatory, Karl-Schwarzschild-Str. 2, 85748 Garching, Germany

⁷⁷ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

⁷⁸ Department of Physics & Astronomy, University College London, Gower Street, London, WC1E 6BT, UK

⁷⁹ Sorbonne Universités, Institut Lagrange de Paris (ILP), 98 bis Boulevard Arago, 75014 Paris, France

⁸⁰ Laboratoire de Physique Nucléaire et de Hautes Energies, Université Pierre et Marie Curie, 4 Place Jussieu, 75005 Paris, France

⁸¹ Shanghai Astronomical Observatory, Chinese Academy of Science, 80 Nandan Road, Shanghai 200030, China

⁸² Department of Astronomy, Case Western Reserve University, Cleveland, OH 44106, USA

⁸³ Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077 Goettingen, Germany

⁸⁴ Academia Sinica Institute of Astronomy and Astrophysics, P.O. Box 23-141, Taipei 10617, Taiwan

⁸⁵ Department of Astronomy, Ohio State University, 140 W. 18th Ave., Columbus, OH 43210, USA

⁸⁶ Center for Cosmology and AstroParticle Physics, The Ohio State University, 191 W. Woodruff Ave., Columbus, OH 43210, USA

⁸⁷ Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85748 Garching, Germany

⁸⁸ Department of Physics, University of Helsinki, Gustaf Hållströmin katu 2a, FI-00014 Helsinki, Finland

⁸⁹ Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA

⁹⁰ Instituto de Astrofísica, Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile

⁹¹ Astrophysical Research Consortium, Physics/Astronomy Building, Rm C319, 3910 15th Avenue NE, Seattle, WA 98195, USA

⁹² National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing 100012, China

⁹³ Tsinghua Center for Astrophysics & Department of Physics, Tsinghua University, Beijing 100084, China

⁹⁴ Astronomical Observatory of Padova, National Institute of Astrophysics, Vicolo Osservatorio 5 - 35122 - Padova, Italy

⁹⁵ Department of Physics, University of North Carolina Asheville, One University Heights, Asheville, NC 28804, USA

⁹⁶ Astrophysics Research Institute, Liverpool John Moores University, IC2, Liverpool Science Park, 146 Brownlow Hill, Liverpool L3 5RF, UK

⁹⁷ SDSS-IV Spokesperson (Corresponding Author)

⁹⁸ Department of Physics and Astronomy, Macquarie University, Sydney NSW 2109, Australia

⁹⁹ ELTE Eötvös Loránd University, Gothard Astrophysical Observatory, Szombathely, Hungary

¹⁰⁰ Premium Postdoctoral Fellow of the Hungarian Academy of Sciences

¹⁰¹ Departamento de Física, Facultad de Ciencias Exactas, Universidad Andres Bello, Av. Fernandez Concha 700, Las Condes, Santiago, Chile.

¹⁰² Vatican Observatory, V00120 Vatican City State, Italy

SDSS-II Supernova Survey (Frieman et al. 2008; Sako et al. 2014), and conducted observations of stars for the Sloan Extension for Galactic Understanding and Exploration 1 (SEGUE-1; Yanny et al. 2009). These surveys made use of the SDSS imaging camera (Gunn et al. 1998) and 640-fiber optical spectrograph (Smee et al. 2013). SDSS-III continued observations of stars with SEGUE-2, and conducted two new surveys with new instrumentation (Eisenstein et al. 2011).

The Baryon Oscillation Spectroscopic Survey (BOSS; Dawson et al. 2013) upgraded the optical spectrograph to 1000 fibers (named the BOSS spectrograph; Smee et al. 2013) to conduct a large volume cosmological redshift survey. The Apache Point Observatory Galactic Evolution Experiment 1 (APOGEE-1; Majewski et al. 2017) employed a high resolution near-infrared spectrograph to observe stars in the Milky Way. All of these observations were conducted at Apache Point Observatory, and data were publicly released in DR12 (Alam, et al. 2015).

This paper contains new data and data reductions produced by SDSS-IV (Blanton, et al. 2017). SDSS-IV began observations in July 2014, and consists of three pro-

grams. The extended Baryon Oscillation Spectroscopic Survey (eBOSS; Dawson et al. 2016) is surveying galaxies and quasars at redshifts $z \sim 0.6\text{--}3.5$ for large scale structure. Mapping Nearby Galaxies at APO (MaNGA; Bundy et al. 2015) is using integral field spectroscopy to study 10,000 nearby galaxies. APOGEE-2 is expanding its observations of the Milky Way; starting in 2017 it began observing the Southern sky, extending its operations to the Las Campanas Observatory’s du Pont Telescope.

Two major subprograms are being conducted concurrently with eBOSS. Spectroscopic IDentification of ERosita Sources (SPIDERS) investigates the nature of X -ray emitting sources, including active galactic nuclei and galaxy clusters. Time Domain Spectroscopic Survey (TDSS; Morganson et al. 2015) is exploring the physical nature of time-variable sources through spectroscopy.

SDSS-IV has had one previous data release (DR13; Albareti et al. 2017; for a “behind the scenes” view of how this is done see Weijmans et al. 2016), which contained the first year of MaNGA data, new calibrations of the SDSS imaging data set, and new processing of APOGEE-1 and BOSS data (along with a small amount of BOSS-related data taken during SDSS-IV).

DR14 contains new reductions and new data for all programs, roughly covering the first two years of SDSS-IV operations. This release contains the first public release of data from eBOSS and APOGEE-2, and almost doubles the number of data cubes publicly available from MaNGA.

The full scope of the data release is described in Section 2, and information on data distribution is given in Section 3. Each of the sub-surveys is described in its own section, with eBOSS (including SPIDERS and TDSS) in Section 4, APOGEE-2 in Section 5, and MaNGA in Section 6. We discuss future plans for SDSS-IV and beyond in Section 7.

2. SCOPE OF DATA RELEASE 14

As has been the case for all public SDSS data releases, DR14 is cumulative, and includes re-releases of all previously released data processed through the most current data reduction pipelines. In some cases this pipeline has not changed for many DR (see summary below). New data released in DR14 were taken by the Sloan Foundation 2.5m telescope between Aug 23 2014 (MJD=56893)¹ and July 10, 2016 (MJD=57580). The full scope of the release is summarized in Table 1.

We discuss the data released by each of the main surveys in detail below, but briefly, DR14 includes:

- Data from 496 new eBOSS plates covering ~ 2480 square degrees observed from September 2014 to May 2016. We also include data from a transitional project between BOSS and eBOSS called the Sloan Extended Quasar, ELG, and LRG Survey (SEQUELS), designed to test target selection algorithms for eBOSS. The complete SEQUELS dataset was previously released in DR13, however DR14 is the first release for eBOSS. The eBOSS data contain mainly Luminous Red Galaxy (LRG) and Quasar spectra, as well as targets from TDSS

¹ this is the date for eBOSS, for APOGEE and MaNGA it was July 2015

¹⁰³ Department of Physics and Astronomy, University of Wyoming, Laramie, WY 82071, USA

¹⁰⁴ University of Alabama, Tuscaloosa, AL 35487, USA

¹⁰⁵ Institut d’Astrophysique de Paris, UMR 7095, CNRS - UPMC, 98bis bd Arago, 75014 Paris, France

¹⁰⁶ Instituto de Física Teórica (IFT) UAM/CSIC, Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain

¹⁰⁷ Instituto de Astrofísica de Andalucía (IAA-CSIC), Glorieta de la Astronomía s/n, E-18008, Granada, Spain

¹⁰⁸ Institut UTINAM, CNRS UMR6213, Univ. Bourgogne Franche-Comté, OSU THETA Franche-Comté-Bourgogne, Observatoire de Besançon, BP 1615, 25010 Besançon Cedex, France

¹⁰⁹ Departamento de Física Teórica M8, Universidad Autónoma de Madrid (UAM), Cantoblanco, E-28049, Madrid, Spain

¹¹⁰ Instituto de Astronomía, Universidad Nacional Autónoma de México, Unidad Académica en Ensenada, Ensenada BC 22860, México

¹¹¹ Department of Physics and Astronomy, Ohio University, Clipping Lab, Athens, OH 45701, USA

¹¹² Institute of Space Sciences (CSIC-IEEC), Carrer de Can Magrans S/N, Campus UAB, Barcelona, E-08193, Spain

¹¹³ Department of Astronomy, University of Illinois, 1002 W. Green Street, Urbana, IL 61801, USA

¹¹⁴ National Center for Supercomputing Applications, 1205 West Clark St., Urbana, IL 61801, USA

¹¹⁵ Stellar Astrophysics Centre, Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, DK-8000 Aarhus C, Denmark

¹¹⁶ Brookhaven National Laboratory, Upton, NY 11973, USA

¹¹⁷ Department of Physical Sciences, The Open University, Milton Keynes, MK7 6AA, UK

¹¹⁸ Kavli Institute for the Physics and Mathematics of the Universe, Todai Institutes for Advanced Study, the University of Tokyo, Kashiwa, Japan 277- 8583

¹¹⁹ Sydney Institute for Astronomy, School of Physics, University of Sydney, NSW 2006, Australia

¹²⁰ School of Physics, University of New South Wales, NSW 2052, Australia

¹²¹ Department of Astronomy and Astrophysics, University of California Santa Cruz, 1156 High St., Santa Cruz, CA, 95064, USA

¹²² Department of Physics, Austin College, Sherman, TX 75090, USA

¹²³ Department of Physics and Astronomy, University of Kentucky, 505 Rose St., Lexington, KY, 40506-0055, USA

¹²⁴ Department of Physics, Bridgewater College, 402 E. College St., Bridgewater, VA 22812 USA

¹²⁵ Hubble Fellow

TABLE 1
REDUCED SPECTROSCOPIC DATA IN DR14

Target Category	# DR13	# DR13+14
eBOSS		
LRG samples	32968	138777
ELG Pilot Survey	14459	35094
Main QSO Sample	33928	188277
Variability Selected QSOs	22756	87270
Other QSO samples	24840	43502
TDSS Targets	17927	57675
SPIDERS Targets	3133	16394
Standard Stars/White Dwarfs	53584	63880
APOGEE-2		
All Stars	164562	263444
NMSU 1-meter stars	894	1018
Telluric stars	17293	27127
APOGEE-N Commissioning stars	11917	12194
MaNGA Cubes	1390	2812
MaNGA main galaxy sample:		
PRIMARY_v1.2	600	1278
SECONDARY_v1.2	473	947
COLOR-ENHANCED_v1.2	216	447
MaNGA ancillary targets ¹	31	121

¹ Many MaNGA ancillary targets were also observed as part of the main galaxy sample, and are counted twice in this table; some ancillary targets are not galaxies.

and SPIDERS. Twenty-three new eBOSS Emission Line Galaxy (ELG) plates are included in DR14 to test final target selection algorithms. The full ELG survey started collecting spectra in September 2016 and will be part of a future data release. We include in DR14 the first part of the ELG target catalogue (see Table 2) described in Raichoor et al. (2017). Other eBOSS value added catalogues (VACs) are also released, namely (1) the redshift measurement and spectral classification catalogue using Redmonster (Hutchinson et al. 2016), (2) the quasar catalogue Paris et al. (in prep.), and (3) a set of composite spectra of quasars binned on spectroscopic parameters (Jensen et al. 2016)).

- APOGEE visit-combined spectra as well as pipeline-derived stellar atmospheric parameters and individual elemental abundances for more than 263,000 stars, sampling all major components of the Milky Way. This release includes all APOGEE-1 data from SDSS-III (Aug 2011-Jul 2014) as well as two years of APOGEE-2 data from SDSS-IV (Jul 2014-Jul 2016). APOGEE VACs include (1) an updated version of the APOGEE red-clump catalog (APOGEE-RC; Pinsonneault et al. in prep.), (2) a cross match between APOGEE and the Tycho-Gaia Astrometric Solution (APOGEE-TGAS; Anders et al. in prep), and (3) a compilation of four different methods to estimate distances to APOGEE stars (Schultheis et al. 2014; Santiago et al. 2016; Wang et al. 2016; Queiroz et al. 2017; Holtzman et al., 2017).
- Data from 166 MaNGA plates, which results in 2812 reconstructed 3D data cubes (for 2744 unique galaxies, primarily from the main MaNGA target sample, but these data also include ancillary targets and ~ 50 repeat observations). Internally this set of galaxies have been referred to as MaNGA

Product Launch-5 (MPL-5); however the reduction pipeline is a different version from that internal release. The new data relative to what was released in DR13 was taken between 13 Aug 2015 (MJD=57248) and July 10 2016. The MaNGA release also includes two VACs, which provide spatially resolved stellar population and ionized gas properties from PIPE3D (Sánchez et al. 2016a,b; see Section 6.4.1) and FIREFLY (Goddard et al. 2017; see Section 6.4.2).

- Overall we include the largest ever number of SDSS Value Added Catalogues (VACs) produced by scientists in the collaboration – twelve in total. See Table 2.
- We re-release the most current reduction of all data from previous versions of SDSS. In some cases the data reduction pipeline hasn’t changed for many DRs, and so has not been re-run. The most recent imaging was released in DR13 (Albareti et al. 2017); however only the photometric calibrations changed in that release; the astrometry is the same as in DR9 (Ahn et al. 2012) and the area released and the other aspects of the photometric reduction remain the same as that in DR8 (Aihara et al. 2011). Legacy Spectra (those observed with the SDSS spectrograph) have also not changed since DR8. There have also been no changes to SEGUE-1 or SEGUE-2 since DR9, or MARVELS since DR12 (Alam, et al. 2015). For DR14 we have re-reduced BOSS spectra using the eBOSS pipeline, where flux calibration has been improved by adding new atmospheric distortion corrections at the per-exposure level (Margala et al. 2016) and by employing an unbiased coaddition algorithm.

3. DATA DISTRIBUTION

TABLE 2
VALUE ADDED CATALOGUES NEW TO DR14

Description	Reference(s)
APOGEE:	
DR14 APOGEE red-clump catalog	Bovy et al. (2014)
DR14 APOGEE-TGAS Catalogue	Anders et al. in prep.
APOGEE DR14 Distance Estimations from Four Groups	
BPG (Bayesian Method)	Santiago et al. (2016); Queiroz et al. (2017)
NAOC (Bayesian Method)	Wang et al. (2016)
NICE (Isochrone Matching Technique)	Schultheis et al. (2014)
NMSU (Bayesian Method)	Holtzman et al., (2017)
eBOSS/TDSS/SPIDERS:	
Redshift Measurement and Spectral Classification Catalog with Redmonster	Hutchinson et al. (2016)
eBOSS: Emission Line Galaxy (ELG) Target Catalog	Raichoor et al. (2017)
The SDSS DR14 Quasar Catalog	Paris et al. in prep.
Composite Spectra of BOSS Quasars Binned on Spectroscopic Parameters from DR12Q	Jensen et al. (2016)
SPIDERS x-ray galaxy cluster catalogue for DR14	Clerc et al. (2016)
Multiwavelength properties of RASS AGN	Merloni et al., in prep
Multiwavelength properties of XMMSL AGN	Del Moro et al., in prep.
MaNGA:	
MaNGA Pipe3D: Spatially resolved and integrated properties of galaxies	Sánchez et al. (2016a,b)
MaNGA FIREFLY Stellar Populations	Goddard et al. (2017)

The DR14 data are distributed through the same mechanisms as DR13, with the addition of a web application to interactively interface with optical and infrared spectra. We describe our three distribution mechanisms below. These methods are also documented on the SDSS website (http://www.sdss.org/dr14/data_access), and tutorial examples for accessing and working with SDSS data can be found at <http://www.sdss.org/dr14/tutorials>.

The raw and processed imaging and spectroscopic data, as well as the value added catalogs, are available through the Science Archive Server (SAS, data.sdss.org/sas/dr14). Data can be downloaded from the SAS directly by browsing the directory structure, and also in bulk using rsync, wget and Globus Online (see http://www.sdss.org/dr14/data_access/bulk for more details). The data files available on the SAS all have their own datamodel, which describes the content of each file in detail. These datamodels are available at <https://data.sdss.org/datamodel>.

The processed imaging and optical and infrared spectra on the SAS are also available through an interactive web application (<https://dr14.sdss.org>). This web application allows the user to search for spectra based on specific parameters, e.g. plate, redshift, coordinates, or observing program. Searches can be saved through permalinks and options are provided to download the spectra directly from the SAS, either individually or in bulk. Previous data releases back to DR8 are available through the same interface. A link is also provided to the SkyServer explore page for each object.

Finally, the DR14 data can be found on the Catalog Archive Server (CAS, Thakar et al. 2008; Thakar 2008). The CAS stores catalogs of photometric, spectroscopic and derived quantities; these are available through the SkyServer web application (<http://skyserver.sdss.org>) for browser-based queries in synchronous mode, and through CasJobs (<http://skyserver.sdss.org/casjobs>), which offers more advanced and extensive query options in asynchronous or batch mode, with more time-consuming queries able to run in the background (Li & Thakar 2008). The CAS is part of the SciServer (<http://www.sciserver.org>) collaborative science framework,

which provides users access to a collection of data-driven collaborative science services, including SkyServer and CasJobs. Other services include SciDrive, a “drag-and-drop” file hosting system that allows users to share files; SkyQuery, a database system for cross-matching astronomical multi-wavelength catalogs; and SciServer Compute, a system that allows users to upload analysis scripts as Jupyter notebooks (supporting Python, MatLab and R) and run these databases in Docker containers.

In addition to the data, the data processing software used by the APOGEE-2, eBOSS, and MaNGA teams to derive their data products from the raw frames, is available at <http://www.sdss.org/dr14/software/products>.

4. EBOSS, TDSS AND SPIDERS

The extended Baryon Oscillation Spectroscopic Survey (eBOSS; Dawson et al. 2016) is surveying galaxies and quasars at redshifts $z \sim 0.6 - 3.5$ to map the large scale structure of the Universe with the main goal to provide Baryonic Acoustic Oscillation (BAO) measurements in the uncharted redshift change spanning $0.6 < z < 2.2$ with a new set of targets: high redshift LRGs, ELGs, and quasars. The three new tracers will provide BAO distance measurements with a precision of 1% at $z = 0.7$ (LRGs), 2% at $z = 0.85$ (ELGs), and 2% at $z = 1.5$ (quasars). The Lyman- α forest imprinted on approximately 120,000 new quasar spectra will give eBOSS an improved BAO measurement of $1.4\times$ over that achieved by BOSS (Delubac et al. 2015; Bautista et al. 2017). Furthermore, the clustering from eBOSS tracers will allow new measurements of redshift-space distortions (RSD), non-Gaussianity in the primordial density field, and the summed mass of neutrino species. eBOSS will provide the first percent-level distance measurements with BAO in the redshift range $0.6 < z < 3$, when cosmic expansion transitioned from deceleration to acceleration. The new redshift coverage of eBOSS obtained by targeting 3 class of targets (LRG, ELG and Quasars) will have the statistical power to improve constraints relative to BOSS by up to a factor of 1.5 in Ω_M , a factor of 3 in the Dark Energy Task Force Figure of Merit (Albrecht et al. 2006),




FIG. 1.— $N(z)$ of eBOSS DR14 QSO and LRG compared to DR12 BOSS, demonstrating how eBOSS is filling in the redshift desert between $z \sim 0.8 - 2.0$.

and a factor of 1.8 in the sum of the neutrino masses (Zhao et al. 2016).

We show in Figure 1 the $N(z)$ in eBOSS DR14 QSO and LRG targets compared to the final BOSS release in DR13 (Alam, et al. 2015), demonstrating how eBOSS is filling in the redshift desert between $z \sim 0.8 - 2.0$.

A significant number of fibers on the eBOSS plates are devoted to two additional dark-time programs. TDSS (Time Domain Spectroscopic Survey; Morganson et al. 2015) seeks to understand the nature of celestial variables by targeting objects that vary in combined SDSS DR9 and Pan-STARRS1 data (PS1; Kaiser et al. 2002). A large number of the likely TDSS quasar targets are also targeted by the main eBOSS algorithms and therefore meet the goals of both surveys. TDSS-only targets fill ~ 10 spectra per square degree. SPIDERS (Spectroscopic Identification of eROSITA Sources) aims to characterize a subset of X-ray sources identified by eROSITA (extended Roentgen Survey with an Imaging Telescope Array; Predehl et al. 2014). However, until the first catalog of eROSITA sources is available, SPIDERS will target sources from the RASS (Roentgen All Sky Survey; Voges et al. 1999) and XMM-Newton (X-ray Multi-mirror Mission; Jansen et al. 2001). SPIDERS will also obtain on average ~ 10 spectra per square degree over the course of SDSS-IV, but the number of fibers per square degree on a plate is weighted toward the later years to take advantage of the new data from eROSITA.

A small fraction of eBOSS time is dedicated to an ancillary program to perform multi-object reverberation mapping for a single 7 deg^2 field. This program (SDSS-RM) aims to detect the lags between the broad-line flux and continuum flux in quasars over a broad range of redshift and luminosity with spectroscopic monitoring, which allows the measurement of the masses of these quasar black holes. Started as an ancillary program in SDSS-III, SDSS-RM continues in SDSS-IV with ~ 12 epochs (each at nominal eBOSS depth) per year to extend the time baseline of the monitoring and to detect lags on multi-year timescales. The details of the SDSS-RM program can be found in Shen et al. (2015), and

initial results on lag detections are reported in Shen et al. (2016) and Grier et al. (2017).

eBOSS started in September 2014 by taking spectra of LRG and Quasars, while further development on the definition of the ELG targets sample was conducted in parallel. In May 2016, eBOSS completed its first major cosmological sample containing LRG and Quasars from the first 2 years eBOSS data and from SEQUELS (already part of the DR13 release). These data have already been used to improve the classification of galaxy spectra (Hutchinson et al. 2016), introduce new techniques to the modeling of incompleteness in galaxy clustering, and to provide measurements of clustering on BAO scales at $1 < z < 2$ for the first time (Ata et al. 2017).

4.1. Data description

DR14 includes the data from 496 plates observed under the eBOSS program; it also includes the 126 SEQUELS plates (already released in DR13), from an ancillary program to take advantage of some of the dark time released when BOSS was completed early. The SEQUELS targets are similar to the eBOSS targets as it was a program to test the selection algorithms of eBOSS, in particular the LRG (Prakash et al. 2016) and quasars (Myers et al. 2015). The final ELG target recipe is not following the one tested during SEQUELS. The new ELG recipe is documented in the DR14 release following the description given by Raichoor et al. (2017).

For the TDSS program, combined SDSS DR9 and Pan-STARRS1 data (PS1; Kaiser et al. 2002) are used to select variable object targets (Morganson et al. 2015; Ruan et al. 2016); while for SPIDERS, the objects are selected from a combination of X-ray and optical imaging for the SPIDERS cluster (Clerc et al. 2016) and AGN (Dwelly et al. 2017) programs.

The sky distribution of the DR14 data from eBOSS is shown in Figure 2. Table 3 summarizes the content and gives brief explanations of the targeting categories.

4.2. Retrieving eBOSS data

All SDSS data releases are cumulative and therefore the eBOSS data also includes the SEQUELS data taken in SDSS-III or SDSS-IV, reduced with the latest pipelines. eBOSS targets can be identified using the EBOSS_TARGET1 bitmask. The summary `spAll-v5.10.0.fits` datafile, which includes classification information from the pipeline, is located on the SAS²; the data can also be queried via the `specObjAll` table on the CAS.

4.3. eBOSS/TDSS/SPIDERS VACs

We include seven VACs based on BOSS, eBOSS, TDSS or SPIDERS data or target selection in this DR. Brief details of each are given below, and for more details we refer you to the relevant papers in Table 2.

4.3.1. Redshift Measurement and Spectral Classification Catalog with Redmonster

The redmonster software³ is a sophisticated and flexible set of Python utilities for redshift measurement, physical parameter measurement, and classification of one-dimensional astronomical spectra. A full description of

² <https://data.sdss.org/sas/dr14/eboss/spectro/redux/v5.10.0/>

³ <https://github.com/timahutchinson/redmonster>




FIG. 2.— DR14 eBOSS spectroscopic coverage in Equatorial coordinates (map centered at RA = 8h.) BOSS coverage is shown in grey, SEQUELS in red, and the eBOSS data newly released for DR14 is shown in blue.

Target Category	Target Flag	# DR14	
Main LRG sample	LRG1_WISE	105764	Prakash et al. (2016)
Ancillary LRG sample	LRG1_IDROP	45	Prakash et al. (2016)
Main QSO selection	QSO1_EBOSS_CORE	154349	Myers et al. (2015)
Variability-selected QSOs	QSO1_VAR_S82	10477	Palanque-Delabrouille et al. (2016)
	QSO1_PTF	54037	Myers et al. (2015)
Reobserved BOSS QSOs	QSO1_REOBS	16333	Myers et al. (2015)
	QSO1_BAD_BOSS	584	
QSOs from FIRST survey	QSO1_EBOSS_FIRST	1792	Myers et al. (2015)
All eBOSS QSOs also in BOSS	QSO_BOSS_TARGET	583	Myers et al. (2015)
All eBOSS QSOs also in SDSS	QSO_SDSS_TARGET	20	Myers et al. (2015)
All “known” QSOs	QSO_KNOWN	11	Myers et al. (2015)
Time-domain spectroscopic survey (TDSS)	TDSS_TARGET	39748	Morganson et al. (2015); MacLeod et al. (2017)
X-ray sources from RASS	SPIDERS_TARGET	13261	Clerc et al. (2016); Dwelly et al. (2017)
X-ray sources in Stripe 82	S82X_TILE1	2775	LaMassa et al in prep. (also see LaMassa et al. 2016)
	S82X_TILE2	2621	
	S82X_TILE3	4	
ELG Pilot Survey	ELG_TEST1	15235	Delubac et al. (2017); Raichoor et al. (2016)
	ELG1_EBOSS	4741	
	ELG1_EXTENDED	659	
Standard stars	STD_FSTAR	8420	Dawson et al. (2016)
Standard white dwarfs	STD_WD	546	Dawson et al. (2016)

the software is given in Hutchinson et al. (2016). The software approaches redshift measurement and classification as a χ^2 minimization problem by cross-correlating the observed spectrum with a theoretically-motivated template within a spectral template class over a discretely sampled redshift interval. In this VAC the software has been run on all DR14 LRG spectra. Redmonster was able to successfully measure redshifts for $\sim 90\%$ of LRG spectra in DR14. This an increase of $\sim 15\%$, in absolute terms, over `spectroid`, and nearly matches the most optimistic estimate for the fraction of measurable redshifts as determined by visual inspections.

4.3.2. The SDSS-IV Extended Baryon Oscillation Spectroscopic Survey: Emission Line Galaxy Target Catalog

We publish the south galactic cap Emission Line Galaxy (ELG) catalog used for eBOSS (Raichoor et al. 2017). Targets were selected using photometric data from the Dark Energy Camera Legacy Survey (DECaLS; <http://legacysurvey.org/>). We selected roughly 240 ELG targets per square degree. The great majority of these ELG lie in the redshift range $0.67 < z < 1.1$ (median redshift 0.85).

4.3.3. The SDSS DR14 Quasar Catalog

Following the tradition established by SDSS-I/II/III the SDSS-IV/eBOSS collaboration is producing a visually inspected quasar catalog. The SDSS-DR14 quasar catalog (DR14Q; Paris et al. in prep.) is the first to be released that contains new identifications that are mostly from eBOSS. The contents of this are similar to the DR12 version (which contained final data from BOSS as well as data from the preliminary eBOSS survey “SEQUELS”) as described in Paris et al. (2017).

4.3.4. Composite Spectra of BOSS Quasars Binned on Spectroscopic Parameters from DR12Q

We present high signal-to-noise composite spectra of quasars over the redshift range $2.1 \leq z \leq 3.5$. These spectra, based on the DR12 BOSS quasar catalog (Alam, et al. 2015) are binned by luminosity, spectral index, and redshift. As discussed in Jensen et al. (2016), these composite spectra can be used to reveal spectral evolution while holding luminosity and spectral index constant. These composite spectra allow investigations into quasar diversity, and can be used to improve the templates used in redshift classification. See Jensen et al. (2017) for more details.

4.3.5. SPIDERS X-ray galaxy cluster catalogue for DR14

A substantial fraction of SPIDERS fibers target red-sequence galaxies in candidate X-ray galaxy clusters. The systems were found by filtering X-ray photon overdensities in the ROSAT All-Sky Survey (RASS) with an optical cluster finder (see Clerc et al. 2016, for details on the samples and targeting strategy). Adding together the DR14 eBOSS sky area with the SEQUELS area (Fig. 1), 573 of these systems show a richness $\lambda_{\text{OPT}} > 30$, have been completely observed as part of DR14 (i.e. all tiled galaxies in a cluster red sequence have got a spectrum in DR14) and contain at least one redshift from SDSS-I to -IV in their red-sequence. Systems located at a border of the DR14 footprint, but in the interior of the full eBOSS footprint, will be fully covered through later observations by overlapping plates.

A total of 9,029 valid redshifts were associated with these candidate rich galaxy clusters, leading to a median number of 15 redshifts per red sequence. An automated algorithm performed a preliminary membership assignment and interloper removal based on standard iterative σ -clipping method. The results of the algorithm were visually inspected by 8 experienced galaxy cluster observers, ensuring at least two independent evaluators per system. A web-based interface was specifically developed to this purpose: using as a starting point the result of the automated algorithm, the tool allows each inspector to interactively assess membership based on high-level diagnostics and figures (see Fig. 16 in Clerc et al. 2016). A final decision is made by each evaluator whether to validate the system as a bona-fide galaxy cluster, or “unvalidate” the system by lack of data or identification of a false candidate. Validation is in most cases a consequence of finding 3 or more red-sequence galaxies in a narrow redshift window, compatible with their being galaxy cluster members. A robust weighted average of the cluster members redshifts provides the cluster systemic redshift. A majority vote was required for each system to be finally “validated” or “unvalidated”; in the former case, an additional condition for agreement is the overlap of the cluster redshifts 95% confidence intervals. A second round of evaluations involving four inspectors per system was necessary to resolve cases with no clear majority.

In total, 520 of these systems are validated as true galaxy clusters based on spectroscopic data and they form the SPIDERS X-ray galaxy cluster Value-Added catalogue for DR14. Among them 478 are unique components along a line-of-sight. A total of 7,352 spectroscopic galaxies are members of a galaxy cluster. This catalogue in particular lists each galaxy cluster redshift and its uncertainty, its number of spectroscopic members and its X-ray luminosity, assuming each component along a line-of-sight contributes the flux measured in RASS data.

4.3.6. Multiwavelength properties of RASS and XMMSL AGN

In these two VACs, we present the multi-wavelength characterization over the area covered by the SEQUELS and eBOSS DR14 surveys (2500 deg^2) of two highly complete samples of X-ray sources; the first from the ROSAT All-Sky Survey (RASS) X-ray source catalogue (2XRS; Boller et al. 2016) and the second from the XMM-

Newton Slew Survey point source catalog (XMMSL; Saxton et al. 2008; version 1.6). We provide information about X-ray properties of the sources, as well as of their counterparts at longer wavelength (Optical, IR, Radio) identified first in the All-WISE IR catalog⁴ via a Bayesian cross-matching algorithm (Dwelly et al. 2017; Salvato et al. 2017, submitted). We complement this with dedicated visual inspection of the SDSS spectra, providing accurate redshift estimates (with objective confidence levels) and source classification, beyond the standard eBOSS pipeline results.

5. APOGEE-2

DR14 is the fourth release from the Apache Point Observatory Galactic Evolution Experiment (APOGEE). DR14 presents, for the first time, the first two years of SDSS-IV APOGEE-2 data (Jul 2014-Jul 2016) as well as re-processed data from SDSS-III APOGEE-1 (Aug 2011-Jul 2014). Note that the general term APOGEE data, employed throughout this paper, refers to both APOGEE-1 and APOGEE-2 data. APOGEE-2 data are substantively the same as APOGEE-1 data, however, one of the three detectors in the instrument was replaced at the end of APOGEE-1 because it exhibited a substantial amount of persistence, in which light from previous exposures leads to excess recorded charge in subsequent exposures; the new detector is substantially better in this regard.

APOGEE data in DR14 includes visit-combined spectra as well as pipeline-derived stellar atmospheric parameters and individual elemental abundances for 263,444 stars⁵, sampling all major components of the Milky Way. In addition to the Milky Way bulge, disk, and halo, DR14 includes, for the first-time, data from stars in satellite galaxies, which are typically fainter targets than those from the main portion of the survey. DR14 incorporates a few modifications in the Data Reduction Pipeline (DRP) as well as in the APOGEE Stellar Parameter and Chemical Abundance Pipeline (ASPCAP). It also includes a separate set of stellar parameters and abundances from The Cannon (Ness et al. 2015)⁶.

Two separate papers will provide more in-depth discussion and analysis of APOGEE data released in DR13/DR14: Holtzman et al. (in prep.) describes in detail the DR13/DR14 pipeline processing as well as the associated data products; and Jönsson et al. (in prep.) compares stellar parameter and element abundances from DR13/DR14 with those from the literature.

5.1. Targeting

The targeting strategy of APOGEE-2 departs slightly from that of APOGEE-1 and is set based on three-tier priority scheme: core, goal, and ancillary (Zasowski et al. in prep.). The core science targets, which are the highest priority, are those that directly address the primary science objectives of APOGEE and include the Galactic bulge, disk, and halo, globular and open clusters, *Kepler* field spectroscopic follow-up, and satellite

⁴ <http://wise2.ipac.caltech.edu/docs/release/allwise/>

⁵ The figure of 263,444 results from the removal of duplicate observations for a single star. Note that DR14 has a total of 277,731 entries.

⁶ Named in recognition of the stellar classification work of Annie-Jump Cannon (Cannon & Pickering 1918).




FIG. 3.— DR14 APOGEE spectroscopic coverage in Galactic coordinates (map centered on the Galactic center). The color coding indicates the number of APOGEE-1+2 targets observed per field, as shown in the key. Fields new to DR14 are outlined in black.

galaxies (unique in APOGEE-2). “Goal” science targets fall in line with APOGEE science goals with a second-tier prioritization and include M dwarfs, eclipsing binaries, substellar companions, *Kepler* Objects of Interest, young (star-forming) clusters, and Extended *Kepler* Mission (K2) spectroscopic follow-up. The third-tier priority are ancillary science targets, for which a general solicitation was issued for programs that could harness the unique capability of the APOGEE instrument. Since the ancillary programs of APOGEE-1 were largely successful and broadened its scientific scope, APOGEE-2 continues in this vein and DR14 presents some of the first ancillary program data. As in APOGEE-1, the primary stellar targets of APOGEE-2 are red giant branch (RGB) stars. APOGEE-2 extends the target stellar classes with designated observations of red clump (RC) stars in the bulge as well as faint stars (e.g., dwarf Spheroidal and halo stream members with $H \geq 14$). On top of the APOGEE-led programs, additional data are collected with the MaNGA co-targeting program. For the MaNGA pointings, APOGEE data is collected concurrently, with the targeted fields in the direction of the Galactic caps. To document the APOGEE-2 targeting scheme, a new set of bit flags is employed in DR14: `APOGEE2.TARGET1`, `APOGEE2.TARGET2`, and `APOGEE2.TARGET3`. Further details with regard to the APOGEE-2 targeting strategy and field design may be found in Zasowski et al. (submitted), including information on APOGEE-2S targets which will be planned to be part of the next data release.

5.2. Pipeline Reduction and Data Product Extraction

As with the previous data releases, all spectra are processed through the DRP, which includes dark current subtraction, cosmic ray removal, flat-fielding, spectral extraction, telluric correction, wavelength calibration, and dither combination. Radial velocities (RVs)

are determined for each individual visit and the individual visit spectra are resampled to rest-wavelength and combined to generate a single spectrum for each object. Associated DRP data products are the visit-combined spectra and radial velocity (RV) values. For DR14, modifications to the RV determination and associated star combination have occurred. The RV values are now determined both relative to the combined spectrum (in an iterative fashion) as well as to the best-matching model. The radial velocities from the method that yields the lower scatter are adopted (`VHELIO_AVG`) and estimates of the associated error and scatter are generated. Note that the new methodology has resulted in improved RV determinations for low signal-to-noise observations (and consequently, faint stars), but there can still be potentially significant issues with some of the faintest targets. The distribution of S/N values for spectra released in DR14 (compared to those released in DR13; Albareti et al. 2017) are shown in Figure 4.

5.2.1. Persistence

As discussed in Nidever et al. (2015) and Holtzman et al. (2015), one of the three APOGEE-1 detectors (the “blue” detector) exhibited significant levels of persistence (i.e., charge which is held between exposures) over one third of the detector area, and another (the “green” detector) exhibited persistence at a somewhat lower level over a smaller area. This persistence affected derived stellar abundances (Holtzman et al. 2015). As mentioned above, the “blue” detector was replaced for APOGEE-2 in part to solve this problem. For the APOGEE-1 data, we attempt to subtract out persistence based on a persistence model and also de-weight pixels affected by persistence during visit combination so that, for stars with a mix of persistence-affected and non-persistence-affected visits, the combined spectra are dominated by the non-




FIG. 4.— A comparison of the S/N distribution of APOGEE spectra released in DR14 (red) with those released in DR13 (blue). The S/N quantity displayed in the figure is SNREV, a revised S/N estimate which considers persistence issues.

affected visits. This results in a reduction in systematic error but a slight increase in random error. This process significantly reduces the impact of persistence (Holtzman et al. in prep); however, it can still have an effect, especially for fainter targets. Users of the APOGEE spectra should pay careful attention to the pixel-level data flags and the pixel uncertainties.

5.2.2. ASPCAP

After the DRP stage, the visit-combined stellar spectra are processed by ASPCAP, which derives the stellar atmospheric parameters (e.g., effective temperature T_{eff} , surface gravity $\log g$, metallicity $[M/H]$) as well as abundances for more than 20 species. The ASPCAP determination proceeds in three stages: an initial pass through ASPCAP gives coarse values for a few key atmospheric parameters to identify which spectral grids should be used on each object, a second pass yields the full set of parameters, and then a final pass determines abundances for each element with stellar parameters fixed. For DR14, ASPCAP modifications include a new normalization scheme for both observed and synthetic spectra. Rather than using an iteratively asymmetrically-clipped fit, the continuum is determined by a polynomial fit to the spectra after masking of sky lines, because the asymmetric clipping is expected to lead to systematic differences in continuum normalization as a function of S/N . Another change is that the ASPCAP parameter determination was done by χ^2 -minimization over a 7-dimensional grid for giants which included a microturbulence dimension. This leads to slightly lower abundance scatter in clusters as well as smaller trends of $[M/H]$ with temperature.

One caveat of the DR14 ASPCAP analysis is that new grids were not constructed for APOGEE-2 line spread functions (LSFs): grids made with the APOGEE-1 LSFs were used. Since the only change was the detector replacement, no large LSF changes were expected, nor were they observed, but the re-installation of the detector package may have led to subtle differences.

5.3. Calibration and Data Product Usage

As with previous DRs, DR14 includes a post-ASPCAP calibration of the final stellar atmospheric parameter and element abundances. A variety of different stellar clusters and standards are employed in the calibration of the results. These calibrations include a metallicity-dependent temperature correction, a surface gravity calibration based on asteroseismic gravities, an internal and external calibration of metallicity ($[M/H]$), and a temperature-dependent and zero-point calibration for elemental abundances. Note that surface gravity calibration is not done for dwarfs because we do not have independent estimates of surface gravities from which to derive such calibrations. Calibrations are applied to abundances over temperature ranges that are determined by looking at the ranges over which data in star clusters produce the same abundance. Based on cluster results and inspection of the spectra, we do not provide calibrated abundances for Cu, Ge, Y, Rb, and Nd, since these do not appear to be reliable.

Several different bitmasks (STARFLAG, PARAMFLAG, ASPCAPFLAG) are included that provide information on factors that affect data quality, and users are strongly encouraged to pay attention to these

5.4. New DR14 Data Product: Results from The Cannon

New in DR14 is the inclusion of parameters and abundances derived from The Cannon (Ness et al. 2015), which has been shown to have the potential to deliver parameters and abundances of higher precision. The Cannon is a data-driven model that provides parameters and abundances (collectively called labels) from the spectra, after training the sensitivity of each pixel to parameters and abundances based on a training set with independently derived labels. For DR14, we train The Cannon on ASPCAP results for a subset of high S/N giant stars, and apply the model to all objects within the range of parameters covered by the training set. DR14 Cannon results have been derived using the Cannon-2 code (Casey et al. 2016), but with a few modifications. First, we adopted uncertainties from the ASPCAP pipeline, which do a better job de-weighting areas around imperfectly subtracted sky lines. Second, and more importantly, we use “censoring” in the derivation of individual elemental abundances, which forces the model to only use pixels where there are known lines of a given element (rather than the full spectrum) to derive the abundance of that element. This was done because it was discovered that, when using the full spectrum, pixels without known lines of an element (and sometimes, with known lines of another element) contributed to the model sensitivity for that element. This suggests that the model may be affected by correlations of abundances within the training set stars. Without censoring, such correlations can lead to abundances that appear to be of higher precision, but this precision may not reflect higher accuracy, if the correlations are not present over the entire data set. While results for some elements with censoring show less scatter than ASPCAP results, results for other elements can look significantly worse. The implementation of censoring was done by using the elemental windows used by the ASPCAP analysis; it is possible that this is overly con-

servative because the ASPCAP windows reject regions in the spectrum that have abundance sensitivity if they are also sensitive to other abundances in the same elemental abundance group.

5.5. APOGEE VACs

Three APOGEE related VACs are included in DR14. They are briefly summarized below. For more details we refer the reader to the relevant paper in Table 2.

5.5.1. DR14 APOGEE red-clump catalog

DR14 contains an updated version of the APOGEE red-clump (APOGEE-RC) catalog. This catalog is created using the same procedure as the original APOGEE-RC catalog (Bovy et al. 2014) now applied to the ASPCAP parameters derived in this data release. To account for changes in how the ASPCAP-derived $\log g$ is calibrated in DR14, we have made the upper $\log g$ cut more stringent by 0.1 dex (the upper $\log g$ limit in Equation [2] in Bovy et al. (2014) now has 2.4 instead of 2.5). Like in the original release, we also apply an additional $\log g$ cut to remove further contaminants (Equation [9] in Bovy et al. 2014). Otherwise the catalog is created in the same manner as the original catalog.

The DR14 APOGEE-RC catalog contains 29,502 unique stars, about 50% more than in DR13. Note that because of changes in the target selection in APOGEE-2, the relative number of RC stars in APOGEE-2 is smaller than in APOGEE-1. We provide proper motions by matching to the UCAC-4 (Zacharias et al. 2013) and HSOY (Altmann et al. 2017) catalogs. Contamination by non-RC stars in the DR14 RC catalog is estimated to be less than 5% by comparing against true RC stars in the APOKASC catalog (M. Pinsonneault et al. in prep.).

5.5.2. DR14 APOGEE-TGAS Catalogue

The first data release of the Gaia mission contains improved parallaxes and proper motions for more than 2 million stars contained in the Tycho-2 catalogue, among them 46,033 objects (10,250 of them unique stars) contained in APOGEE DR14. This is known as the Tycho-Gaia Astrometric Solution (TGAS). We provide the cross-matched catalog, together with precise combined astrometric/spectro-photometric distances and extinctions determined with *StarHorse* (Queiroz et al. 2017) for 29,661 stars. We also include orbital parameters calculated using the *GravPot16* code⁷ (Fernandez-Trincado et al., in prep.). For more details see Anders et al. (in prep.).

5.5.3. APOGEE DR14 Distance Estimations from Four Groups

This VAC provides spectro-photometric distance estimates for APOGEE stars that have been calculated by four groups, using slightly different isochrone techniques. All groups used the DR14 calibrated ASPCAP stellar parameters, if they fall inside the calibration ranges (see Holtzman et al., in prep.). The distances come from (1) the *StarHorse* code (Santiago et al. 2016; Queiroz et al. 2017, in prep.) (2) the code described in Wang et al. (2016), (3) the isochrone-matching technique described in Schultheis et al. (2014), and (4) the distance code described in Holtzman et al., (2017).

⁷ <https://fernandez-trincado.github.io/GravPot16/>

6. MANGA

In the context of the MaNGA Survey, DR14 roughly doubles the sample size of associated data products that were first made public in DR13. Spanning observations from the first two years of operations, the DR14 products include raw observations, intermediate reduction output, such as reduced fiber spectra, and final data cubes as constructed by the Data Analysis Pipeline (Law et al. 2016, , L16 hereafter). A summary *drpall* catalog provides target identification information, sky positions, and object properties like photometry and redshifts. The MaNGA observing strategy is described in Law et al. (2015), and the flux calibration scheme presented in Yan et al. (2016b). An overview of the survey execution strategy and data quality is provided in Yan et al. (2016a). Weijmans (2016) provides a short summary to the entire survey, which is comprehensively described in Bundy et al. (2015).

DR14 includes observations from 166 MaNGA plates resulting in 2812 datacubes comprising targets in the main samples as well as ancillary programmes, and around 50 repeat observations. The sky layout of the DR14 released MaNGA data is shown in Figure 5.

6.1. MaNGA Target Classes

The target selection for the MaNGA Survey is described in detail by Wake et al. (2017). MaNGA’s main galaxy sample is complete to stellar masses, $M_* > 10^9 M_\odot$, and is comprised of three main sub-samples that are defined on the basis of SDSS-I/II photometry and spectroscopic redshifts to deliver a final distribution that is roughly flat in $\log M_*$. The Primary sample achieves radial coverage out to 1.5 times the effective radii ($1.5 R_e$) for target galaxies, while the Secondary sample reaches $2.5 R_e$. The Color-Enhanced supplement expands the selection of the Primary sample to include under-represented regions of M_* -color space. We refer to the combination of the Primary and Color-Enhanced supplements as “Primary+” which balances the rest-frame color distribution at fixed M_* . The MaNGA samples can be weighted so that they are equivalent to a volume limited sample. The required volume weights are described in Wake et al. (in prep.) and are provided in the DR14 version of the targeting file.

DR14 includes 1278 Primary galaxies, 947 Secondary galaxies and 447 Color-Enhanced supplement galaxies. Which sample a given target galaxy belongs to is given by the `MANGA_TARGET1` bitmask (or `mngtarg1` in the “*drpall*” file). Bits 10, 11, and 12 signal that galaxies were selected as Primary, Secondary, or Color-Enhanced targets respectively. In addition to ~ 121 ancillary program targets, ~ 50 galaxies were observed as fillers and do not fall into these target categories. They should be ignored in statistical studies of the MaNGA data.

MaNGA has also begun observing Milky Way stars in a bright-time survey program called the MaNGA Stellar Library (MaStar) that makes use of MaNGA IFUs during APOGEE-2 observations. The goal of MaStar is to build a new stellar library comprising > 8000 stars that span the widest accessible ranges in effective temperature, surface gravity, metallicity, and element abundance ratios (Yan et al. in preparation). Reduced stellar spectra will be included in DR15.




FIG. 5.— The sky distribution (Mollweide equatorial projection for Dec $> -20^\circ$) of possible MaNGA plates (in grey). Because MaNGA targets are selected from a sample with SDSS-I photometry and redshifts, this footprint corresponds to the Data Release 7 imaging data (Abazajian et al. 2009). Each plate contains 17 MaNGA targets, and around 30% of all possible plates will be observed in the full 6-year survey. The purple indicates plates with data released as part of DR14.

As described in the DR13 paper (Albareti et al. 2017, hereafter DR13), roughly 5% of MaNGA IFUs are allocated to targets defined by approved ancillary programs. These sources can be identified using the MANGA_TARGET3 bitmask (or mngtarg3 in the drpall file). Most of the programs represented in DR14 are described in DR13⁸. They include targeted followup of AGN hosts, starburst galaxies, merging systems, dwarf galaxies, Milky Way analogs, and brightest cluster galaxies. New in DR14, we include deep observations reaching ~ 20 hours in the center of the Coma cluster (Gu et al. in prep.) and IFU observations allocated as part of an ancillary program to a nearby dwarf galaxy that is part of the ACS Nearby Galaxy Survey (Dalcanton et al. 2009).

6.2. Working with MaNGA Data

All MaNGA data products take the form of multi-extension FITS files. As we describe in DR13, the DRP data products consist of intermediate reduced data (sky-subtracted, flux-calibrated fiber spectra with red and blue data combined for individual exposures of a plate) and final-stage data products (summary row-stacked spectra and data cubes) for each target galaxy. The summary row stacked spectra (RSS files) are two-dimensional arrays provided for each galaxy in which each row corresponds to a single fiber spectrum.

The three-dimensional data cubes are created by combining the individual spectra for a given galaxy together onto a regularized $0.5''$ grid (see L16 for more detail). The associated wavelength arrays for both the data cubes and RSS files can be accessed in logarithmic and linear scales. Each data cube contains additional extensions with information that includes the inverse variance, a bad-pixel mask, instrumental resolution, reconstructed broadband images, and the effective spatial point-spread function. The full data model for all MaNGA DRP data products can be found online at <http://www.sdss.org/dr14/manga/manga-data/data-model> and in Appendix B of L16.

Instructions for accessing the MaNGA data products are given on the SDSS website⁹. We summarize available options here and refer the reader to the DR13 paper for

additional details. All data products are stored on the Science Archive Server at <http://data.sdss.org/sas/dr14/manga/spectro/redux/> which includes the drpall summary table and subdirectories that store reduction output for each plate, both for observations obtained on a specific night and for the results of combining all observations of a given plate into a “stack.” The drpall table may be queried either after downloading this file to disk or through the SDSS CASJobs system. Such queries define selections of galaxies of interest and can return the plate-IFU combination for those galaxies that identifies how they were observed. These in turn can be used to find the SAS directory locations of the corresponding data products. Large downloads can be accomplished via rsync calls as described on the SDSS website. Finally, the SDSS SkyServer Explore tool provides basic information about MaNGA targets.

Several features of the MaNGA data should be kept in mind while using the data. Most important, each MaNGA data cube has a FITS header keyword DRP3QUAL indicating the quality of the reduction. 1-2% of the data cubes are flagged as significantly problematic—galaxies with CRITICAL quality bit (=30) set should be treated with extreme caution (see L16). Please also use the MASK extension of each datacube to identify problematic spaxels. A simple summary DONOTUSE bit is of particular importance indicating elements that should be masked out for scientific analyses.

There is significant covariance between adjacent spaxels in data cubes, given that the spaxel size ($0.5''$) is much smaller than the fiber size ($2''$ diameter). A simple method that accounts for covariance when one desires to spatially bin spaxels together is discussed in §9.3 of L16. The typical reconstructed point spread function of the MaNGA data cubes has a FWHM of $2.5''$. Sparse correlation matrices in the *ugriz* central wavelengths are also now provided in the data cubes.

As discussed by L16, the instrumental line spread function (LSF) in the DR13 data was underestimated by about $10 \pm 2\%$. This has been corrected in DR14 and the reported LSF is described by a post-pixellized gaussian.

Additional issues and caveats are discussed in <http://www.sdss.org/dr14/manga/manga-caveats/>.

6.3. Highlights of MaNGA Science with DR14 Data

⁸ also see <http://www.sdss.org/dr14/manga/manga-target-selection/ancillary-targets>

⁹ <http://www.sdss.org/dr14/manga/manga-data/data-access/>

The MaNGA survey has produced a number of scientific results based on data acquired so far, indicating the breadth of research possible with the MaNGA data. In Albareti et al (2017) we provide a summary of science highlights with early data. Here we briefly summarize the results of papers that have been completed within the SDSS-IV collaboration using the MaNGA sample released as part of DR14.

For example, published results based on the MaNGA DR14 data include, Barrera-Ballesteros et al. (2017) who discuss the integrated stellar mass-metallicity relation for more than 1700 galaxies, Zhu et al. (2017) who revisit the relation between the stellar surface density, the gas surface density and the gas-phase metallicity of typical disc galaxies in the local Universe, Belfiore et al. (2017) who study the gas phase metallicity and nitrogen abundance gradients traced by star-forming regions in a representative sample of 550 nearby galaxies, and Lin et al. (2017) who report the discovery of a mysterious giant H α blob that is ~ 8 kpc away from a component of a dry galaxy merger. In Bizyaev et al. (2017) was presented a study of the kinematics of the extraplanar ionized gas around several dozen galaxies, while Jones et al. (2017) conducted a detailed study of extra-planar diffuse ionized gas stacking spectra from 49 edge-on, late-type galaxies as a function of distance from the midplane of the galaxy. Numerous other results based on DR14 data are in preparation.

6.4. MaNGA VACs

This data release also contains two VACs based on MaNGA data. They are briefly summarized below, and for more details we refer you to the papers given in Table 2

6.4.1. MaNGA Pipe3D value added catalog: Spatially resolved and integrated properties of galaxies

PIPE3D is an IFU-focused analysis pipeline that calculates intermediate dataproducts and is able to obtain both the stellar population and the ionized gas properties extracted from the datacubes in an automatic way. This pipeline is based on FIT3D, details of which are presented in Sánchez et al. (2016a,b) which show some examples based on CALIFA (Sánchez-Menguiano et al. 2016; Cano-Díaz et al. 2016; Sánchez et al. 2017) and MaNGA/P-MaNGA (Ibarra-Medel et al. 2016; Barrera-Ballesteros et al. 2016; Lin et al. 2017; Barrera-Ballesteros et al. 2017) datasets. The MaNGA dataproducts provided by Pipe3D will be presented in Sánchez et al. (in prep)¹⁰. The VAC consists of a single table containing integrated (cumulative), characteristic (values at the effective radius), and gradient of different quantities, included stellar mass, star-formation (and their densities), oxygen and nitrogen abundances, dust attenuation, estimated gas density, stellar and gas velocity dispersions.

For each galaxy, data is presented as an individual FITS file including four extensions, each one corresponding to a data cube that comprises (1) the spatial resolved properties required to recover the star-formation histories, (2) the average properties of the stellar populations, (3) the emission line properties for 56 strong and weak

emission lines (including the former ones together with the EW of the lines), and (4) the most frequently used stellar indices. The details of each individual extension was described in Sánchez et al. (2016b), and the final adopted format in Sánchez et al. (in prep.).

6.4.2. MaNGA FIREFLY Stellar Populations

The MaNGA FIREFLY VAC (Goddard et al. 2017) provides measurements of spatially resolved stellar population properties in MaNGA galaxies. It is built on and complements the products of the MaNGA data analysis pipeline (DAP, Westfall et al, in preparation) by providing higher-order and model-based data products. These are measurements of optical absorption line-strengths, as well as the physical properties age, metallicity and dust attenuation. The latter are derived from full spectral fitting with the code FIREFLY (Wilkinson et al. 2015, 2017) using the supercomputer SCIAM2 at Portsmouth University. The VAC is a single fits file (4 GB) containing measurements all DR14 MaNGA galaxies. The catalogue contains basic galaxy information from the literature (i.e., galaxy identifiers, redshift, mass), global derived parameters (i.e., light-weighted and mass-weighted stellar population ages and metallicities for a central 3 arcsec aperture and for an elliptical shell at 1 effective radius), gradient parameters (i.e., gradients in age and metallicity measured within $1.5 R_e$) and spatially resolved quantities (i.e., 2-D maps of age, metallicity, dust attenuation, mass and surface mass density, and 28 absorption line indices).

More detail on the catalogue, and the method of creating the 2-dimensional maps is provided in Goddard et al. (2017) and the data is available from the data release website¹¹.

7. FUTURE PLANS

SDSS-IV is planning a 6-year survey, with operations at both the 2.5 meter Sloan Foundation Telescope at Apache Point Observatory, New Mexico, USA and the du Pont Telescope at Las Campanas, Chile scheduled through 2020. Future data releases from SDSS-IV will include data observed with both telescopes; the final SDSS-IV data release is planned to be DR18, currently scheduled for December 2020.

For APOGEE, future data releases will include, for the first time, southern hemisphere observations taken with the new APOGEE-S instrument at the Las Campanas Observatory with the duPont 2.5m telescope. These observations will extend APOGEE coverage to the full Galaxy, with significantly increased observations of the Galactic bulge and also include observations in the Magellanic Clouds, globular clusters, and dwarf spheroidal galaxies only accessible from the southern hemisphere. As usual, future data releases will also include re-reductions of all APOGEE-N data. Plans for improved stellar parameter/abundance analysis include using a new homogeneous grid of MARCS stellar atmospheres, and the use of “minigrids” to analyze elements whose absorption features are too blended with those of other elements to be reliably extracted with the abundance techniques used to date.

¹⁰ <http://www.sdss.org/dr14/manga/manga-data/manga-pipe3d-value-added-catalog>

¹¹ <http://www.sdss.org/dr14/manga/manga-data/manga-firefly-value-added-catalog>

For MaNGA, it is planned that the DR15 data release will include the ~ 4000 MaNGA galaxies which have been observed up to summer shutdown 2017. In addition, we anticipate a number of new data products to be released in this and future DRs. These include reduced spectra from the MaStar stellar library (Yan et al. in prep.), which is making use of commensal observations during APOGEE-2 time to obtain spectroscopic observations of stars which will be used to build a new stellar library through the MaNGA instrumentation; and output from the MaNGA Data Analysis Pipeline (DAP; Westfall et al. in prep.). The DAP produces maps of emission line fluxes, gas and stellar kinematics, and stellar population properties. Some similar derived data products are already available as Value Added Catalogs (see Table 2 and Section 6.4.1 and 6.4.2). Finally, we intend for DR15 to mark the first release of the “Marvin” ecosystem which includes powerful python tools for seamlessly downloading and querying the MaNGA data as well as web interface that provides advanced search functionality, an user interface to the MaNGA datacubes, and the ability to quickly choose and display maps of key quantities measured by the DAP.

For eBOSS, future data releases will include the ELG survey results as well as the continuation of the LRG-QSO surveys. They will also include further value added catalogues: in particular the continuation of the quasar catalogue, a detailed ELG catalogue, as well as large scale structure clustering catalogues required for independent clustering analysis. Further improvement on the redshift measurement and spectral classification catalogue is also likely.

For TDSS, a future SDSS data release will also include very recent spectra from its Repeat Quasar Spectroscopy (RQS) program, which obtains multi-epoch spectra for thousands of known quasars, already with at least one epoch of SDSS spectroscopy available (and often already archived). Quasar spectral variability on multi-year timescales is heretofore relatively poorly characterized for large samples, but there are of course many exciting results from smaller select subsets (see Runnoe et al. 2016 and McGraw et al. 2017 for examples of studies based on repeat spectroscopy, ranging from discoveries of new changing look quasars, to BAL emergence and disappearance). Observations for the RQS program in TDSS, which will ultimately observe $\sim 10^4$ known (SDSS) quasars in the ELG survey region (Raichoor et al. 2017) for at least one additional spectral epoch in SDSS-IV, extend such earlier work to a more systematic investigation of quasar spectroscopic variability; this extension is via both larger samples, but also by including large numbers of quasars as targets for repeat spectra that were selected without marked *a priori* biases based on specific quasar spectroscopic sub-class or variability character. A recent detailed technical description of target selection for all of the TDSS repeat spectroscopy programs (including RQS), may be found in MacLeod et al.

(2017); and RQS spectra will be included in a future SDSS data-release.

For SPIDERS, future data releases will focus on higher-level data products, such as black hole masses and host galaxy properties of the X-ray AGN, as well as rich characterization of the X-ray selected clusters (in particular, dynamical properties and calibrated cluster masses). The first spectra of counterparts of eROSITA sources, however, will only be obtained beginning in Spring 2019, so they will be part of DR18 and subsequent releases only.

Planning has begun for the After SDSS-IV (AS4) project, to begin in 2020. AS4 will build on the SDSS infrastructure and expand the instrumentation (especially for optical IFU spectroscopy) in both hemispheres. This expansion of SDSS’s legacy will enable an enormous sample of millions of spectra of quasars, galaxies, and stars, with scientific goals ranging from the growth of super-massive black holes to the detailed architecture of planetary systems and the astrophysics of stellar interiors.

8. ACKNOWLEDGEMENTS

We would like to thank the University of St Andrews, Scotland for their hospitality during DocuCeilidh 2017.

Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS-IV acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS web site is www.sdss.org.

SDSS-IV is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, the Chilean Participation Group, the French Participation Group, Harvard-Smithsonian Center for Astrophysics, Instituto de Astrofísica de Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU) / University of Tokyo, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), Max-Planck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatories of China, New Mexico State University, New York University, University of Notre Dame, Observatório Nacional / MCTI, The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, University of Washington, University of Wisconsin, Vanderbilt University, and Yale University.

REFERENCES

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 543
- Albrecht, A., Bernstein, G., Cahn, R., et al. 2006, *ArXiv Astrophysics e-prints*
- Ahn, C. P., Alexandroff, R., Allende Prieto, C., et al. 2012, *ApJS*, 203, 21
- Aihara, H., et al. 2011, *ApJS*, 193, 29
- Alam, S. et al. 2015, *ApJS*, 219, 12

- Albareti, F. D., Allende Prieto, C., et al. 2017, *ApJS* submitted (arXiv:1608.02013)
- Altmann, M., Roeser, S., Demleitner, M., Bastian, U., & Schilbach, E. 2017, *A&A*, 600, L4
- Ata, M., Baumgarten, F., Bautista, J., et al. 2017, arXiv:1705.06373
- Barrera-Ballesteros, J. K., Heckman, T. M., Zhu, G. B., et al. 2016, *MNRAS*, 463, 2513
- Barrera-Ballesteros, J. K., Sánchez, S. F., Heckman, T., & Blanc, G. A. 2017, *ApJ* in press (arXiv:1706.09893)
- Bautista, J. E., Busca, N. G., Guy, J., et al. 2017, arXiv:1702.00176
- Belfiore, F., Maiolino, R., Tremonti, C., et al. 2017, *MNRAS*, 469, 151
- Bizyaev, D., Walterbos, R. A. M., Yoachim, P., et al. 2017, *ApJ*, 839, 87
- Blanton, M. R., Bershad, M. A., Abolfathi, B., et al. 2017, *AJ*, 154, 28
- Boller, T., Freyberg, M. J., Trümper, J., et al. 2016, *A&A*, 588, A103
- Bovy, J., Nidever, D. L., Rix, H.-W., et al. 2014, *ApJ*, 790, 127
- Bundy, K., et al. 2015, *ApJ*, 798, 7
- Cannon, A. J., & Pickering, E. C. 1918, *Annals of Harvard College Observatory*, 91, 1
- Cano-Díaz, M., Sánchez, S. F., Zibetti, S., et al. 2016, *ApJ*, 821, L26
- Casey, A. R., Hogg, D. W., Ness, M., et al. 2016, arXiv:1603.03040
- Chevallier, M., & Polarski, D. 2001, *International Journal of Modern Physics D*, 10, 213
- Comparat, J., Delubac, T., Jouvel, S., et al. 2016, *A&A*, 592, A121
- Clerc, N., Merloni, A., Zhang, Y.-Y., et al. 2016, *MNRAS*, 463, 4490
- Dalcanton, J. J., Williams, B. F., Seth, A. C., et al. 2009, *ApJS*, 183, 67
- Dawson, K. S., Schlegel, D. J., Ahn, C. P., et al. 2013, *AJ*, 145, 10
- Dawson, K., et al. 2016, *AJ*, 151, 44
- Delubac, T., Bautista, J. E., Busca, N. G., et al. 2015, *A&A*, 574, A59
- Delubac, T., Raichoor, A., Comparat, J., et al. 2017, *MNRAS*, 465, 1831
- Dwelly, T., Salvato, M., Merloni, A., et al. 2017, *MNRAS*, 469, 1065
- Eisenstein, D. J., Weinberg, D. H., Agol, E., et al. 2011, *AJ*, 142, 72
- Frieman, J. A., Bassett, B., Becker, A., et al. 2008, *AJ*, 135, 338
- Goddard, D., Thomas, D., Maraston, C., et al. 2017, *MNRAS*, 466, 4731
- Grier, C. J., Trump, J. R., Shen, Y., et al. 2017, *ApJ*, submitted
- Gunn, J. E., Carr, M., Rockosi, C., et al. 1998, *AJ*, 116, 3040
- Gunn et al. 2006, *AJ* 131, 2332
- Holtzman, J. A., Shetrone, M., Johnson, J. A., et al. 2015, *AJ*, 150, 148
- Holtzman et al., in prep
- Hutchinson, T. A., Bolton, A. S., Dawson, K. S., et al. 2016, *AJ*, 152, 205
- Ibarra-Medel, H. J., Sánchez, S. F., Avila-Reese, V., et al. 2016, *MNRAS*, 463, 2799
- Jansen, F., Lumb, D., Altieri, B., et al. 2001, *A&A*, 365, L1
- Jensen, T. W., Vivek, M., Dawson, K. S., et al. 2016, *ApJ*, 833, 199
- Jones, A., Kauffmann, G., D'Souza, R., et al. 2017, *A&A*, 599, A141
- Kaiser, N., Aussel, H., Burke, B. E., et al. 2002, *Proc. SPIE*, 4836, 154
- LaMassa, S. M., Urry, C. M., Cappelluti, N., et al. 2016, *ApJ*, 817, 172
- Law, D. R., Yan, R., Bershad, M. A., et al. 2015, *AJ*, 150, 19
- Law, D. R., Cherinka, B., Yan, R., et al. 2016, *AJ*, 152, 83
- Li, N., & Thakar, A. R. 2008, *Computing in Science and Engineering*, 10, 18
- Lin, L., Lin, J.-H., Hsu, C.-H., et al. 2017, *ApJ*, 837, 32
- Linder, E., & SNAP Collaboration 2002, *Bulletin of the American Astronomical Society*, 34, 97.02
- MacLeod, C. L., Green, P. J., Anderson, S. F., et al. 2017, arXiv:1706.04240
- Madrid, J. P., & Macchetto, D. 2009, *BAAS*, 41, 913
- Majewski, S. R., Schiavon, R. P., Frinchaboy, P. M., et al. 2017, *AJ*, submitted, arXiv:1509.05420
- Margala, D., Kirkby, D., Dawson, K., et al. 2016, *ApJ*, 831, 157
- McGraw, S. M., Brandt, W. N., Grier, C. J., et al. 2017, *MNRAS*, 469, 3163
- Morganson, E., Green, P. J., Anderson, S. F., et al. 2015, *ApJ*, 806, 244
- Myers, A. D., Palanque-Delabrouille, N., Prakash, A., et al. 2015, *ApJS*, 221, 27
- Nidever, D. L., Holtzman, J. A., Allende Prieto, C., et al. 2015, *AJ*, 150, 173
- Palanque-Delabrouille, N., Magneville, C., Yèche, C., et al. 2016, *A&A*, 587, A41
- Pâris, I., Petitjean, P., Ross, N. P., et al. 2017, *A&A*, 597, A79
- Prakash, A., Licquia, T. C., Newman, J. A., et al. 2016, *ApJS*, 224, 34
- Predehl, P., Andritschke, R., Becker, W., et al. 2014, *Proc. SPIE*, 9144, 91441T
- Queiroz et al. 2017 in prep.
- Raddick, M. Jordan, Ani R. Thakar, Alexander S. Szalay, & Rafael DC Santos. 2014, *Computing in Science and Engineering* 16, 22.
- Raddick, M. Jordan, Ani R. Thakar, Alexander S. Szalay, & Rafael DC Santos. 2014, *Computing in Science and Engineering* 16, 32.
- Raichoor, A., Comparat, J., Delubac, T., et al. 2016, *A&A*, 585, A50
- Raichoor, A., Comparat, J., Delubac, T., et al. 2017, *MNRAS* submitted (arXiv:1704.00338)
- Ruan, J. J., Anderson, S. F., Cales, S. L., et al. 2016, *ApJ*, 826, 188
- Runnøe, J. C., Cales, S., Ruan, J. J., et al. 2016, *MNRAS*, 455, 1691
- Sako, M., Bassett, B., Becker, A. C., et al. 2014, arXiv:1401.3317
- Sánchez, S. F., Pérez, E., Sánchez-Blázquez, P., et al. 2016a, *RMxAA*, 52, 21
- Sánchez, S. F., Pérez, E., Sánchez-Blázquez, P., et al. 2016b, *RMxAA*, 52, 171
- Sánchez, S. F., Barrera-Ballesteros, J. K., Sánchez-Menguiano, L., et al. 2017, *MNRAS*, 469, 2121
- Sánchez-Menguiano, L., Sánchez, S. F., Pérez, I., et al. 2016, *A&A*, 587, A70
- Santiago, B. X., Brauer, D. E., Anders, F., et al. 2016, *A&A*, 585, A42
- Saxton, R. D., Read, A. M., Esquej, P., et al. 2008, *A&A*, 480, 611
- Schultheis, M., Zasowski, G., Allende Prieto, C., et al. 2014, *AJ*, 148, 24
- Shen, Y., Brandt, W. N., Dawson, K. S., et al. 2015, *ApJS*, 216, 4
- Shen, Y., Horne, K., Grier, C. J., et al. 2016, *ApJ*, 818, 30
- Smee, S. A., et al. 2013, *AJ*, 146, 32
- Stoughton, C., Lupton, R. H., Bernardi, M., et al. 2002, *AJ*, 123, 485
- Thakar, A. R., Szalay, A., Fekete, G., & Gray, J. 2008, *Computing in Science and Engineering*, 10, 30
- Thakar, A. R. 2008, *Computing in Science and Engineering*, 10, 9
- Voges, W., Aschenbach, B., Boller, T., et al. 1999, *A&A*, 349, 389
- Wake, D. A., Bundy, K., Diamond-Stanic, A. M., et al. 2017, arXiv:1707.02989
- Wang, J., Shi, J., Pan, K., et al. 2016, *MNRAS*, 460, 3179
- Weijmans, A.-M., & MaNGA Team 2016, *Multi-Object Spectroscopy in the Next Decade: Big Questions, Large Surveys, and Wide Fields*, 507, 257 (arXiv:1508.04314)
- Weijmans, A.-M., Blanton, M., Bolton, A. S., et al. 2016, arXiv:1612.05668
- Wilkinson, D. M., Maraston, C., Thomas, D., et al. 2015, *MNRAS*, 449, 328
- Wilkinson, D. M., et al. 2017, *MNRAS*, in press.
- Wylezalek, D., Schnorr Müller, A., Zakamska, N. L., et al. 2017, *MNRAS*, 467, 2612
- Yan, R., Bundy, K., Law, D. R., et al. 2016a, *AJ*, 152, 197
- Yan, R., Tremonti, C., Bershad, M. A., et al. 2016b, *AJ*, 151, 8
- Yanny, B., Rockosi, C., Newberg, H. J., et al. 2009, *AJ*, 137, 4377-4399
- York, D. G., et al. 2000, *AJ*, 120, 1579
- Zacharias, N., Finch, C. T., Girard, T. M., et al. 2013, *AJ*, 145, 44

Zhao, G.-B., Wang, Y., Ross, A. J., et al. 2016, MNRAS, 457,
2377
Zhu, G. B., Barrera-Ballesteros, J. K., Heckman, T. M., et al.
2017, MNRAS, 468, 4494